



**UNIVERSITI PUTRA MALAYSIA**

**INDUCED RESISTANCE USING NONPATHOGENIC FUSARIUM  
OXYSPORUM FOR BIOLOGICAL CONTROL OF BANANA  
FUSARIUM WILT**

**TARIG ELSAYED ALI ELSHARIF**

**FP 2003 32**

**INDUCED RESISTANCE USING NONPATHOGENIC *FUSARIUM*  
*OXYSPORUM* FOR BIOLOGICAL CONTROL OF BANANA *FUSARIUM*  
WILT**

**By**

**TARIG ELSAYED ALI ELSHARIF**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirement for the Degree of Doctor of Philosophy**

**February 2003**



## **DEDICATION**

**I dedicate this humble effort, the fruit of my thoughts and study to my affectionate Mother, Father, wife Nagwa and children who inspired me to higher ideals of life.**

Abstract of thesis submitted to the Senate of Universiti Putra Malaysia in fulfilment of the requirements for the degree of Doctor of Philosophy

**INDUCED RESISTANCE USING NONPATHOGENIC *FUSARIUM*  
*OXYSPORUM* FOR BIOLOGICAL CONTROL OF BANANA *FUSARIUM*  
WILT**

By

**TARIG ELSAYED ALI ELSHARIF**

**February 2003**

**Chairperson : Professor Dr. Sariah Meon**

**Faculty : Agriculture**

Banana (*Musa* spp. Linn.) is the second most important fruit crops in Malaysia. It is easily attacked by the fungus *Fusarium oxysporum* Schlecht. f.sp. *cubense* (E.F.Smith) Snyder and Hansen (FOC), causing terminal wilt in the field. Chemical and cultural methods were not effective in controlling the disease. Therefore, alternative control measures have to be developed. In recent years, nonpathogenic *Fusaria* was being considered for plant disease control and could be the most promising approach for biological control of *Fusarium* wilt in banana through induced systemic resistance.

Isolates of nonpathogenic *F. oxysporum* (FO: FO1, FO2, FO3, FO4, FO5 and FO6) were isolated from healthy roots and rhizospheres of bananas vars. *Berangan* and *Rastali*, and were identified to the species level based on cultural and morphological characteristics. Random amplified polymorphic (RAPD-PCR) analysis was able to



establish variability within *F. oxysporum* isolates and between saprophytic and pathogenic forms (FOC race 1 and race 4), using OPC 11 and OPC 14 primers. All 6 isolates of FO were antagonistic to both pathogenic race 1 and race 4 of FOC with values of the % of inhibition of radial growth (PIRG) exceeding 50% in a series of dual culture test. FO4 was found to be the most antagonistic against FOC race 4 with PIRG of 65%.

Infectivity studies on six-weeks-old tissue cultured banana seedlings var. *Berangan* cv. *Intan*, confirmed that FO1, FO2, FO3, FO4, FO5 or FO6 were not pathogenic to banana seedlings. No visible foliar or internal symptoms were observed both on inoculated and control seedlings. Seedlings inoculated with FOC race 4 produced foliar symptoms as yellowing of the older leaves followed by necrosis and wilting. FO4 conferred some degree of resistance to the host when challenged with FOC race 1 suggesting the possible role of induced resistance against *Fusarium* wilt.

In this experiment, it was shown that resistance of banana seedlings var. *Berangan* cv. *Intan* against FOC race 4 was associated with inducible compounds such as peroxidases (PO) and polyphenol oxidases (PPO). There was a sharp increase in total PO and PPO activities in the early phase of infection (3-12 days) before the appearance of foliar symptoms. In the case of FOC race 4 infected activities of PO and PPO were lowest at the end of the assessment, and correspond with development of severe foliar and internal symptoms. Production of gum-like substances and formation of tyloses in xylem vessels were apparent in FO4 inoculated seedlings. Hypersensitive reactions (HR) were evident which resulted in localized infection. PO and PPO were associated with the deposition of lignin, an

important component of cell wall, chemically modified to be more resistant to cellular degrading enzymes. The proline accumulation in the leaves of FOC race 4 inoculated plants was significantly higher (at  $p \geq 0.05$ ) when compared to those of FO4 inoculated plants and control plants throughout the duration of the experiment, suggesting that proline could be used as a marker of stress.

#### Tissue-cultured

and nine days with inducer FO4 and later challenged with FOC race 4. In all treatments FO4 able to control *Fusarium* wilt disease. Disease severity (%) as marked by values of Area Under the Disease Progress Curve (AUDPC) was significantly lower compared to single inoculation with FOC race 4. An interval of nine days between inducer and challenger inoculation gave 21.8% reduction in disease severity after 54 days of challenged inoculation. Lignin accumulation in the roots was 1.2 ug/g FW as compared to 0.9 ug/g FW in non-treated seedlings based on the lignin thioglycolic acid (LTGA) assay.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**KERESISTANAN TERARUH MENGGUNAKAN *FUSARIUM OXYSPORUM*  
BUKAN PATOGENIK UNTUK KAWALAN BIOLOGI LAYU FUSARIUM  
PISANG**

Oleh

**TARIG ELSAYED ALI ELSHARIF**

**February 2003**

**Pengerusi : Professor Dr. Sariah Meon**

**Fakulti : Pertanian**

Pisang (*Musa* spp. Linn.) adalah tanaman buah-buahan yang penting di Malaysia. Ianya mudah di serang oleh kulat *Fusarium oxysporum* Schlecht. f.sp. *cubense* (E.F.Smith) Snyder and Hansen (FOC), mengakibatkan penyakit layu vaskular di ladang. Kawalan kimia dan kultur tidak berkesan untuk mengawal penyakit ini. Oleh yang demikian kawalan alternatif perlu diusahakan. Beberapa tahun kebelakangan ini, Fusaria bukan patogenik telah dipertimbangkan untuk pegawalan penyakit tumbuhan dan mungkin merupakan pendekatan kawalan biologi terbaik untuk mengawal penyakit layu fusarium pisang melalui aruhan resistan sistemik.

*F. oxysporum* bukan pathogenic (FO1, FO2, FO3, FO4, FO5 dan FO6) telah di pencilkan daripada akar dan rhizosfera pokok pisang var. *Berangan* dan *Rastali* yang sakit dan dikenalpasti sehingga peringkat sepsis berdasarkan ciri kultur dan morfologi. Analisis "Random amplified polymorphic" (RAPD-PCR) membolehkan pengecaman terperinci diantara isolat *F. oxysporum* dan di antara bentuk saprofitik dan patogenik (FOC ras 1 dan FOC ras 4), menggunakan primer- primer OPC 11 dan

OPC 14. Kesemua isolat FO ini adalah antagonistik terhadap FOC (ras 1 dan-ras 4) dengan peratusan perencatan pertumbuhan radial (PIRG) melebihi 50% dalam siri ujian dwikultur. FO4 telah didapati lebih antagonistik terhadap FOC race 4 dengan PIRG 65%.

Kajian kesan jangkitan ke atas anak benih kultur tisu pisang *Berangan* cv. *Intan* berumur enam minggu mengesahkan FO1, FO2, FO3, FO4, FO5 atau FO6 adalah bukan patogenik terhadap anak benih pisang. Tiada simptom pada daun atau simptom dalaman yang dapat dilihat Pada kedua-dua anak benih yang diinokulat dan kawalan. Anak benih yang dijangkiti oleh FOC ras 4 menunjukkan simptom foliar dengan daun tua menjadi kuning diikuti oleh nekrosis dan layu. FO4 dapat meningkatkan pertumbuhan vegetatif anak benih pisang dan memberi darjah kerekesitanan pada perumah bila dicabar dengan FOC ras 1 mencadangkan kemungkinan peranan kerekesitanan raruhan terhadap layu Fusarium pada anak benih yang diuji.

Dalam kajian ini menunjukkan Keresistanan anak benih pisang var. *Berangan* cv. *Intan* terhadap FOC ras 4 dikaitkan dengan kompaun perangsang, peroxidase (PO) dan polifenoloxidase (PPO). Terdapat peningkatan yang amat ketara dalam jumlah PO dan PPO di awal jangkitan (3 hingga 12 hari) sebelum kemunculan symptom foliar. Pada pokok yang dijangkiti oleh FOC ras 4, aktiviti PO dan PPO adalah sangat rendah diakhir kajian dan dikaitkan dengan pembentukan symptom foliar dan dalaman yang teruk. Penghasilan bahan seperti gam dan pembentukan tylosis dalam saluran xilem adalah jelas pada anak benih jangkitan FO4. Reaksi hipersensitif (HR), adalah jelas dan menyebabkan infeksi setempat. PO dan PPO berkait rapat dengan

penendapan lignin, komponen penting dalam dinding sel, mengubahsuai dinding sel secara kimia menjadikan ia lebih resistan terhadap enzim pereputan selular. Pengumpulan proline dalam daun yang dijangkiti FOC ras 4 adalah lebih tinggi dan bererti (pada  $P \geq 0.05$ ) dibandingkan dengan FO4 dan kawalan sepanjang tempoh ujikaji, mencadangkan proline boleh digunakan sebagai penanda bagi tegasan.

Anak benih pisang tisu kultur var *Berangan* cv. *Intan* yang di pra-jangkitkan selama tiga, enam dan sembilan hari dengan FO4 dan kemudian dicabar dengan FOC ras 4. Dalam semua rawatan FO4 mampu mengawal penyakit layu *Fusarium*. Peratus keterukan penyakit yang digambarkan oleh kawasan di bawah keluk perkembangan penyakit (AUDPC) adalah rendah dibandingkan dengan anak benih yang dijangkitkan hanya dengan FOC ras 4 tempoh sembilan hari di antara jangkitan perangsang dan pencabar memberikan 21.8% penurunan keterukan penyakit 54 hari selepas jangkitan cabaran. Pengumpulan lignin dalam akar ialah 1.2 ug/g berbanding dengan 0.9 ug/g dalam anak benih tanpa rawatan berdasarkan esei penentuan asid lignin thioglycolic (LTGA).

## ACKNOWLEDGMENTS

All praises and thanks are of Allah, the beneficent and merciful. The author invokes Allah's blessing peace for the Holy Prophet Muhammad (peace be upon him). The messenger of Allah, who advised all of us to continue getting education from cradle till death.

I avail myself of the opportunity in recording my sincerest thanks and appreciations to Professor Dr. Sariah Meon, the head of supervisory committee, Professor Dr. Marziah Mahmud and Associate Professor Dr. Kamaruzzaman Sijam as members of the supervisory committee, for their guidance, understanding, constructive criticism, suggestions and invaluable advice throughout the duration of this study and preparation of this thesis.

I am exceedingly grateful to Sudan Government, Omdurman Islamic University (Sudan), Sudan Embassy in KL, Malaysia Government and MTCP for their financial support and without their help the present project would have been a mere dream.

Also with deep sense of honour I wish to extend my sincere gratitude to Universiti Putra Malaysia, Graduate school and Faculty of Agriculture for their assistance that make my study successful and to United Plantations Berhad for supplying the tissue-cultured plant materials used in this study. Thanks are also due to Professor Dr. Yusof Hussein former Dean Faculty of Agriculture for his financial support.

Thanks are also extended to all staffs in the Pathology and Nematology Laboratory Mr. Khir, Mr. Nazri, Mr. Johari, Mr. Lim K.A, Mr. Gani, and Mr. Shamsuddin for their advice and assistance. Special thanks to Ms. Afaf M, Prof. Hassan Elnagar, Mr Babikir, Dr. Yosif M., Elwaleed A., Ismail, Chow, Hendry and Eltahir S.

I take this opportunity to express my deepest gratitude to my mother, brothers, sisters and my wife Nagwa and daughters Elaf, Linah, Alaa, and sons Muhammad and Elsayed Ali who exhibited great patience, goodwill, encouragement, love and understanding throughout the period of my study.

I certify that an Examination Committee met on 5<sup>th</sup> February 2003 to conduct the final Examination of Tarig Elsyed Ali El Sharif on His Doctor of Philosophy thesis entitled "Induced Resistance using Nonpathogenic *Fusarium oxysporum* for Biological Control of Banana Fusarium Wilt" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Zainal Abidin Mior Ahmad, Ph.D.**

Lecturer,  
Faculty of Agriculture,  
Universiti Putra Malaysia.  
(Chairman)

**Sariah Meon, Ph.D.**

Professor,  
Faculty of Agriculture,  
Universiti Putra Malaysia.  
(Member)

**Kamaruzaman Sijam, Ph.D.**

Associate Professor,  
Faculty of Agriculture,  
Universiti Putra Malaysia.  
(Member)

**Marziah Mahmood, Ph.D.**

Professor,  
Faculty of Science and Environmental Studies,  
Universiti Putra Malaysia.  
(Member)

**Baharuddin Salleh, Ph.D.**

Professor,  
School of Biological Sciences,  
Universiti Sains Malaysia.  
(Independent Examiner)



---

**SHAMSHER MOHAMAD RAMADILI, Ph.D.**

Professor/Deputy Dean,  
School of Graduate Studies,  
Universiti Putra Malaysia.

Date: . 5 MAR 2003



This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:

**Sariah Moon, Ph.D.**

Professor,  
Faculty of Agriculture,  
Universiti Putra Malaysia.  
(Chairman)

**Kamaruzaman Sijam, Ph.D.**

Associate Professor,  
Faculty of Agriculture,  
Universiti Putra Malaysia.  
(Member)

**Marziah Mahmood, Ph.D.**

Professor,  
Faculty of Science and Environmental Studies,  
Universiti Putra Malaysia.  
(Member)

---

**AINI IDERIS, Ph.D.**

Professor/Dean,  
School of Graduate Studies,  
Universiti Putra Malaysia.

Date:

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



---

**TARIG ELSAYED ALI ELSHARIF**

Date: **27.02.03**

## TABLE OF CONTENTS

DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVALS	xi
DECLARATION	xiii
LIST OF TABLE	xvii
LIST OF FIGURES	xviii
LIST OF PLATES	xx
LIST OF ABBREVIATIONS	xxii

## CHAPTER

1	INTRODUCTION	1.1
1.1	Objectives of Study	1.6
1.2	Hypothesis	1.6
2	LITERATURE REVIEW	2.1
2.1	Systematic and Characterization of <i>Fusarium oxysporum</i>	2.1
2.2	Characterization of <i>Fusarium oxysporum</i> f. sp. <i>cubense</i>	2.2
2.3	Fusarium Wilt of Banana	2.5
2.4	Disease Distribution and Losses	2.8
2.5	Epidemiology	2.10
2.6	Physiological Studies on Pathogenesis	2.12
2.6.1	Proline	2.12
2.6.2	Nett Photosynthesis	2.12
2.6.3	Chlorophylls	2.13
2.6.4	Stomatal Conductance	2.13
2.7	Management of Fusarium Wilt Disease	2.14
2.7.1	Cultural and Physical Control	2.14
2.7.2	Chemical Control	2.15
2.7.3	Biological Control	2.17
2.7.3.1	Suppressive Soil	2.17
2.7.3.2	Microbial Antagonists	2.19
2.8	Induced Resistance Using Nonpathogenic Strains	2.20
2.8.1	Mechanisms of Induced Resistance	2.24
2.8.1.1	Glucanases and Chitinases	2.26
2.8.1.2	Peroxidases	2.26
2.8.1.3	Polyphenol Oxidases	2.28
2.8.1.4	Phenolic Compounds	2.29
2.8.1.5	Physical Barriers	2.30



3	ISOLATION AND CHARACTERISATION OF NONPATHOGENIC STRAINS OF <i>FUSARIUM</i> <i>OXYSPORUM</i>	3.1
3.1	Introduction	3.1
3.2	Materials and Methods	3.3
3.2.1	Isolation of <i>Fusarium</i> Isolates	3.3
3.2.2	Characterisation of <i>Fusarium</i> Isolates	3.4
3.2.2.1	Morphological and Cultural Characteristics	3.4
3.2.2.2	DNA- Polymorphism	3.4
3.2.2.3	<i>In vitro</i> Antagonistic Activities of <i>Fusarium</i> Isolates	3.7
3.2.3	Effects of <i>Fusarium</i> Isolates on Growth and Disease Severity of Banana Seedlings	3.9
3.2.3.1	Infectivity Studies	3.9
3.2.3.2	Disease Suppression	3.11
3.3	Results and Discussion	3.13
3.3.1	Isolation and Characterisation of <i>Fusarium</i> isolates	3.13
3.3.1.1	RAPD-PCR Amplification	3.17
3.3.1.2	<i>In- vitro</i> Antagonistic Activities of <i>F.</i> <i>oxysporum</i> Isolates	3.22
3.3.2	Effect of <i>F. oxysporum</i> Isolates on Growth and Disease Severity of Banana Seedlings	3.25
3.3.2.1	Infectivity Studies	3.25
3.3.2.2	Disease Suppression	3.35
4	MECHANISMS OF INDUCED RESISTANCE AGAINST FUSARIUM WILT	4.1
4.1	Introduction	4.1
4.2	Materials and Methods	4.3
4.2.1	Growth Responses	4.4
4.2.2	Disease Development	4.4
4.2.3	Physiological Responses	4.4
4.2.3.1	Stomatal Conductance	4.4
4.2.3.2	Nett Photosynthesis	4.4
4.2.3.3	Chlorophyll Content	4.5
4.4.4	Biochemical Responses	4.5
4.4.4.1	Protein	4.6
4.4.4.2	Peroxidase and Polyphenol Oxidase	4.6
4.4.4.3	Proline	4.8
4.4.5	Histological Responses	4.9
4.3	Results and Discussion	4.12

4.3.1	Growth Responses	4.12
4.3.2	Development of Fusarium Wilt	4.14
4.3.3	Physiological Responses	4.19
4.3.4	Biochemical Responses	4.25
	4.3.4.1 Peroxidase (PO)	4.25
	4.3.4.2 Polyphenol Oxidase (PPO)	4.31
4.3.5	Proline	4.33
4.3.6	Histological Responses	4.36
5	EFFECTIVENESS OF FO4 AS AN INDUCER OF SYSTEMIC RESISTANCE IN BANANA SEEDLINGS	5.1
5.1	Introduction	5.1
5.2	Materials and Methods	5.1
	5.2.1 Lignin Assay	5.3
5.3	Results and Discussion	5.5
	5.3.1 Disease Severity	5.5
	5.3.2 Plant Growth	5.14
	5.3.3 Lignin	5.16
6	GENERAL DISCUSSION AND CONCLUSION	6.1
	REFERENCES	R.1
	APPENDICES	A.1
	BIODATA OF THE AUTHOR	B.1

## LIST OF TABLES

Table		Page
3.1	Frequency of isolation of fungi from banana rhizosphere var. <i>Rastali</i> and <i>Berangan</i> on <i>Fusarium</i> semi selective media	3.14
3.2	Mean (n=6) <i>in vitro</i> antagonistic activities of <i>F. oxysporum</i> isolates against FOC race 1 and FOC race 4.	3.24
3.3	Mean (n=6) disease severity index (foliar symptoms) of <i>Fusarium</i> wilt on banana seedling var. <i>Berangan</i> cv. <i>Intan</i> inoculated with FOC race1 and race 4.	3.31
3.4	Mean (n=6) Area Under Disease Progress Curve (AUDPC) of <i>Fusarium</i> wilt in banana seedlings pre-inoculated with nonpathogenic <i>F. oxysporum</i> and challenged with FOC race 1.	3.38
4.1	Mean (n=6) disease severity index value of FOC race 4 inoculated seedlings according to days.	4.16
5.1	Treatments (n=6) carried out in evaluating FO4 as an inducer of resistance in banana seedlings var <i>Berangan</i> cv <i>Intan</i> challenged with FOC race 4.	5.2
5.2	Mean (n=6) Area under disease progress curve (AUDPC) of <i>Fusarium</i> wilt in banana seedlings challenged with FOC race 4, at 3, 6, 9 days after pre-inoculated with FO4.	5.8
5.3	Mean (n=6) disease severity (foliar) index value (DSI) of banana seedlings var <i>Berangan</i> cv. <i>Intan</i> pre-inoculated with FO4 and challenged with FOC race 4.	5.9
5.4	Mean (n=6) disease severity (vascular discoloration of corm) index (DSI) of banana seedlings var <i>Berangan</i> cv. <i>Intan</i> challenged with FOC race 4 at 54 days pre-inculcated with FO4.	5.9
5.5	Mean (n=6) lignin concentration in root tissues and disease severity index of banana seedlings var. <i>Berangan</i> cv. <i>Intan</i> challenged with FOC race 4, 3 days (FR1), 6 days (FR2) and 9 (FR3) days after pre-inoculated with FO4, values were determined 54 days after inoculation.	5.18

## LIST OF FIGURES

Figure		Page
3.1	Phenogram Based on RAPD-PCR Profiles Generated by UPGMA Analysis Showing the Genetic Relationships Among Pathogenic Isolates of FOC race 1 and race 4 and Nonpathogenic Isolates (FO1, FO2, FO3, FO4, FO5 and FO6) with Primers OPC 11 and OPC 14.	3.19
3.2	Mean (n=5) Progress of Fusarium Wilt on Banana Seedling var. <i>Berangan</i> cv. <i>Intan</i> Inoculated with FOC Race1 (R1) and Race 4 (R4). Bars Represent the Standard Deviation of the Difference Between Means.	3.30
3.3	Mean (n=6) Disease Severity % in var <i>Berangan</i> cv <i>Intan</i> after Six Weeks of Inoculation with FOC Race 1 (R1) & FOC Race 4 (R4). Bars Represent the Standard Deviation of the Difference Between Means.	3.31
3.4	Mean (n=6) Increase in Plant Height and Stem Diameter in var. <i>Berangan</i> cv. <i>Intan</i> Inoculated with Pathogenic and Nonpathogenic Isolates of <i>Fusarium</i> , Six Weeks after Inoculation.	3.33
3.5	Mean (n=6) Increase in Leaf Length of var <i>Berangan</i> cv. <i>Intan</i> Inoculated with Pathogenic and Nonpathogenic Isolates of <i>Fusarium</i> , Six Weeks after Inoculation.	3.34
3.6	Mean (n=6) Disease Severity (%) in Banana Seedlings var. <i>Berangan</i> cv. <i>Intan</i> After Six Weeks of Re-Inoculation With FOC race 1.	3.37
4.1	Mean (n=6) Increase in Height (cm) of Banana Seedlings var <i>Berangan</i> cv <i>Intan</i> Inoculated with FO4 and FOC race 4. Bars Represent the Standard Deviation of the Difference Between Means.	4.13
4.2	Mean (n=6) Increase in Stem Diameter (mm) of Banana Seedlings var <i>Berangan</i> cv. <i>Intan</i> Inoculated with FO4 and FOC race 4. Bars Represent the Standard Deviation of the Difference Between Means.	4.13
4.3	Mean (n=6) Development of Foliar Symptoms in Banana Seedlings var <i>Berangan</i> cv. <i>Intan</i> Inoculated with <i>F. oxysporum</i> f.sp.cubense race 4. Bars Represent the Standard Deviation of the Difference Between Means.	4.15
4.4	Mean (n=6) Nett Photosynthesis ( $\text{mmol m}^{-2} \text{ s}^{-1}$ ) and Stomatal Conductance ( $\text{mmol m}^{-2} \text{ s}^{-1}$ ) of Banana Seedlings var.	

	<i>Berangan</i> cv <i>Intan</i> Inoculated with <i>F. oxysporum</i> (FO4) and <i>F. oxysporum</i> f.sp. <i>cubense</i> race 4 (FOCR4); In Relation to Disease Severity. Bars Represent the Standard Deviation of the Difference Between Means.	4.22
4.5	Mean (n=6) Chlorophyll Content (mg/cm <sup>2</sup> ) on Leaf of Banana Seedlings var. <i>Berangan</i> cv. <i>Intan</i> Inoculated with FO4 and FOC race 4; in Relation to Disease Severity. Bars Represent the Standard Deviation of the Difference Between Means.	4.24
4.6	Mean (n=6) Total Peroxidase (PO) Activity in Leaf Tissues of Banana Seedlings var. <i>Berangan</i> cv. <i>Intan</i> Inoculated with <i>F.oxysporum</i> (FO4) and <i>F.oxysporum</i> f.sp. <i>cubense</i> race 4; In Relation to Disease Severity. PO Activity is Expressed as Change Absorbance per ug Protein at 470nm. Bars Represent the Standard Deviation of the Difference Between Means.	4.27
4.7	Mean (n=6) Total Peroxidase (PO) Activity in Root Tissues of Banana Seedlings var <i>Berangan</i> cv <i>Intan</i> Inoculated with <i>F.oxysporum</i> (FO4) and <i>F.oxysporum</i> f.sp. <i>cubense</i> race 4; In Relation to Disease Severity. PO Activity is Expressed as Change in Absorbance per ug Protein at 470nm. Bars Represent the Standard Deviation of the Difference Between Means.	4.29
4.8	Mean (n=6) Proline Concentration (ug/g) in Leaf Tissues of Banana Seedlings var <i>Berangan</i> cv <i>Intan</i> Inoculated with <i>F.oxysporum</i> (FO4) and <i>F.oxysporum</i> f.sp. <i>cubense</i> race 4, In Relation to Disease Severity. (Arrows indicate DSI values). Bars Represent the Standard Deviation of the Difference Between Means.	4.35
5.1	Mean (n=6) Disease Progress of Fusarium Wilt on Banana Seedlings var. <i>Berangan</i> cv. <i>Intan</i> Challenged with FOC race 4 at 3, 6 and 9 Days after Pre-inoculated with Inducer (FO4). Bars Represent the Standard Deviation of the Difference Between Means.	5.7
5.2	Mean (n=6) Increase in Plant Height (PH) and Stem Diameter (SD) of Banana Seedlings var. <i>Berangan</i> cv. <i>Intan</i> , 54 Days after Challenged with FOC race 4.	5.15
5.3	Lignin (n=6) Concentration in Root Tissues of Banana Seedlings var. <i>Berangan</i> cv. <i>Intan</i> Challenged with FOC race 4, 3 Days (FR1), 6 Days (FR2) and 9 (FR3) Days after Pre-inoculation with FO4, Values were Determined 54 Days after Inoculation. Relative Lignin Content was Determined by the Thioglycolic Acid Procedure and Expressed as ug/ mg F.W.	5.17



## LIST OF PLATES

Plate		Page
3.1	Morphological Characteristics of <i>Fusarium oxysporum</i> (FO4) Showing; Globose to Falcate Macroconidia (A), Chlamydospores (B) and Microconidia (C).	3.16
3.2	RAPD of Nonpathogenic Isolates of <i>F. oxysporum</i> Obtained with Primers OPC 11 and OPC 14. Lane 1 to 6 Contain Isolates of <i>F. oxysporum</i> (FO1, FO2, FO3, FO4, FO5 and FO6), Isolated from Roots and Rhizosphere of Banana var. <i>Rastali</i> and <i>Berangan</i> , Lane 7 to 8 Contain Pathogenic Isolates of FOC race 4 and race 1, M Contain DNA Marker (bp).	3.18
3.3	Antagonistic Activities of Six Isolates of <i>Fusarium oxysporum</i> (FO1-FO6) Against FOC race 1 and FOC race 4 Based in Dual Culture Test.	3.23
3.4	Disease Severity Scale for the Assessment of Foliar Symptoms on Banana Seedlings var. <i>Berangan</i> cv. <i>Intan</i> .	3.26
3.5	Disease Severity Scale for the Assessment of Internal Symptoms of the Corm on Banana Seedlings var <i>Berangan</i> cv <i>Intan</i> .	3.27
3.6	Internal Symptoms in the Root (A), Stem (B) and Corm (C) of Banana Seedlings Infected by FOC race 4 (arrow). Co: Control.	3.29
3.7	Development of Foliar Symptoms on Banana Seedlings var <i>Berangan</i> cv <i>Intan</i> . Challenged with <i>F.oxysporum</i> f. sp. <i>cubense</i> race 1, Six Weeks after Pre-Inoculation with Inducer <i>F. oxysporum</i> (FO4).	3.36
3.8	Internal Symptoms on Corm of Banana Seedlings var. <i>Berangan</i> Inoculated with FOC race 1 (left), Pre-inoculated with FO4 and Challenged with <i>F.oxysporum</i> f. sp. <i>cubense</i> race 1 (right).	3.39
4.1A	Progress of Internal Symptoms in Root and Corm (vascular discoloration) of Banana Seedlings var <i>Berangan</i> cv <i>Intan</i> . Inoculated with FOC race 4 (as indicated by arrow).	4.18
4.1B	Hypersensitive Reactions (HR) Caused by FO4 on Banana Root (as indicated by arrow).	4.18

4.2	Longitudinal Section on Banana Root Inoculated with FOC race 4, Showing Mycelium (M), Gummosis (G) and Tyloses (T) in Xylem Vessel.	4.37
4.3	Scanning Electron Micrograph on Banana Root var. <i>Berangan</i> cv. <i>Intan</i> Inoculated with FOC race 4, Showing Tyloses Emerging (A) from Pits in Scalariform Wall of Infected Vessel. Infected Tyloses (B).	4.38
4.4	Cross Section of Banana Root var. <i>Berangan</i> cv. <i>Intan</i> , Showing Deposition of Lignin (thickness) in the Wall of the Cells Around Xylem Vessel, (Arrow), after Inoculation with to <i>F. oxysporum</i> (FO4) and <i>F. oxysporum</i> f.sp. <i>cubense</i> race 4 (FOCR4). C: control.	4.39

## LIST OF ABBREVIATIONS

µg	Microgram
ANOVA	Analysis of variance
AUDPC	Area Under Disease Progress Curve
bp	base pair
CRD	Complete Randomised Design
cfu	Colony forming units
cm	centimetre
dd	double distilled water
DMRT	Duncan's multiple range test
DNA	Deoxy ribonucleic acid
DS	Disease severity
DSI	Disease severity index
EDTA	Ethylene diaminetetra acetic acid
FA	Fusaric acid
FAO	Food and Agriculture Organization
FO	<i>Fusarium oxysporum</i>
FOC	<i>Fusarium oxysporum</i> f.sp. <i>cubense</i>
FW	fresh weight
g	Gram
ha	hectare
h	Hour
HCL	Hydrochloric acid
HR	Hypersensitive
IS	Induced resistance
ISR	Induced Systemic Resistance
Kb	Kilo-base pair
Kg	Kilogram (10 <sup>3</sup> gram)
L	litre
LTGA	ligninthioglycolic acid
M	Molar
MARDI	Malaysia Agriculture Research and Development Institute
Min	Minute
mL	Mililiter
mm	Milimeter
mM	Milimolar
Mmolm <sup>-2</sup> s <sup>-1</sup>	milimole per meter square per second
NaCl	Sodium chloride
NaOH	Sodium hydroxide.
ng	Nanogram (10 <sup>-9</sup> microgram)
nm	Nanometer
NPK	Nitrogen, Phosphorous, Potassium
°C	Centigrade
OPC	Oligo-nucleotide purification column primers
<i>p</i>	probability
PCR	Polymerase chain reaction

PIRG	percentage of inhibition of radial growth
PO	Peroxidase
PPO	polyphenol oxidase
RAPD	Random Amplified Polymorphic DNA
SAR	Systemic acquired resistance
SC	Similarity coefficient
spp.	species
TAE	Tris –HCL-glacial acetic acid-EDTA
TE	Tris-EDTA
ug	microgram ( $10^{-3}$ gram)
ul	microliter ( $10^{-3}$ ml)
UPM	Universiti Putra Malaysia
UV	Ultra violet
v/v	volume/volume
VCGs	Vegetative C complete ompatibility Groups

## CHAPTER I

### INTRODUCTION

Banana (*Musa* spp. (Linn.)) belonging to the Family Musaceae (Wardlaw, 1972), is the second most important fruit crop in Malaysia, accounting for 20% of the total hectareage for planted fruit (Doon, 1991; Siti Hawa, 2000). The banana industry has the potential to become one of the most profiting fruit commodities in future, marked by the increase in demand and consumption, that is expected to generate an increase in production to over 400 000 ha. by the year 2010 (Rohizad, 1998). However, land used for banana cultivation has declined over the years from 40 000 ha. in 1993 to 39 000 ha. in 2000 (Loh, 2000). Traditionally, banana cultivation is confined to smallholdings as a cash crop or as a temporary intercrop with oil palm or other crops. The popular dessert cultivars grown for domestic market are *Pisang Mas* (AA), *Pisang Embun* (AAA), *Pisang Rastali* (AAB) and *Pisang Berangan* (AAA). The cooking cultivars are *Pisang Tanduk* (AAB), *Pisang-Raja* (AAB), *Pisang Nangka* (AAB), *Pisang Awak* (ABB) and *Pisang Abu* (ABB). *Pisang Mas* is the most important cultivar grown for domestic consumption and for export, followed by *Pisang Berangan*, *Pisang Embun* and *Pisang Rastali*.

Both sweet and starchy cooking types of banana called plantains, are staple food of most tropical countries. In addition, bananas and plantains provide high sources of dietary carbohydrate, vitamin C, and a number of important minerals (Tezenas, 1991). Total world production is estimated at over 76 million metric tons (FAO, 1993), of which exports (essentially of *Cavendish* bananas) to the richer nations represent less than 11 million tons. World banana production is nearly 50 million tons of which over 40% is in Asia.